

AD-A193 193

ENANTIOSELECTIVE RING-CLEAVAGE OF MESO- EPOXIDES WITH
BETA-HALODIISOPINOCAMPHEYLBORANES(U) PURDUE UNIV
LAFAYETTE IN N N JOSHI ET AL. 26 APR 88 TR-1
N00014-86-K-0547


1/1

UNCLASSIFIED

F/G 7/3

NL





MICROCOPY RESOLUTION TEST CHART

10101 10101 10101 10101 10101

4

OFFICE OF NAVAL RESEARCH

DTIC FILE COPY

Contract N00014-86-K-0547

R&T Code 4135011---02

Technical Report No. 1

Enantioselective Ring-cleavage of *meso*- Epoxides
with B-Halodiisopinocampheylboranes

by

N. N. Joshi, M. Srebnik, and Herbert C. Brown

Prepared for Publication

in the

Journal of American Chemical Society

Purdue University
H. C. Brown and R. B. Wetherill Laboratories of Chemistry
West Lafayette, Indiana 47907

April 26, 1988

DTIC
ELECTE
MAY 03 1988
S D
H

Reproduction in whole or in part is permitted for
any purpose of the United States Government.

* This document has been approved for public release
and sale; its distribution is unlimited.

AD-A193 193

88 5_02 308

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

| | | | | | |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|----------------------------|
| 1a. REPORT SECURITY CLASSIFICATION Unclassified | | | 1b. RESTRICTIVE MARKINGS | | |
| 2a. SECURITY CLASSIFICATION AUTHORITY | | | 3. DISTRIBUTION / AVAILABILITY OF REPORT Distribution List Enclosed | | |
| 2b. DECLASSIFICATION / DOWNGRADING SCHEDULE | | | | | |
| 4. PERFORMING ORGANIZATION REPORT NUMBER(S) | | | 5. MONITORING ORGANIZATION REPORT NUMBER(S) | | |
| 6a. NAME OF PERFORMING ORGANIZATION Purdue University | | 6b. OFFICE SYMBOL (If applicable) | | 7a. NAME OF MONITORING ORGANIZATION Office of Naval Research | |
| 6c. ADDRESS (City, State, and ZIP Code) West Lafayette, IN 47907 | | | 7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217 | | |
| 8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research | | 8b. OFFICE SYMBOL (If applicable) | | 9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0547 | |
| 8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217-5000 | | 10. SOURCE OF FUNDING NUMBERS | | | |
| | | PROGRAM ELEMENT NO. | PROJECT NO. | TASK NO. | WORK UNIT ACCESSION NO. |
| 11. TITLE (Include Security Classification) Enantioselective Ring Cleavage of <i>meso</i> -Epoxides with <i>R</i> -Halodiisopinocampheylboranes | | | | | |
| 12. PERSONAL AUTHOR(S) N. N. Joshi, M. Srebnik and H. C. Brown | | | | | |
| 13a. TYPE OF REPORT Technical | | 13b. TIME COVERED FROM _____ TO _____ | | 14. DATE OF REPORT (Year, Month, Day) April 26, 1988 | |
| 15. PAGE COUNT 12 | | | | | |
| 16. SUPPLEMENTARY NOTATION | | | | | |
| 17. COSATI CODES | | | 18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Ipc ₂ BX, <i>meso</i> -epoxides, enantioselective ring cleavage, 1,2-halohydrins | | |
| FIELD | GROUP | SUB-GROUP | | | |
| | | | | | |
| | | | | | |
| 19. ABSTRACT (Continue on reverse if necessary and identify by block number) see attached report | | | | | |
| 20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS | | | 21. ABSTRACT SECURITY CLASSIFICATION Unclassified | | |
| 22a. NAME OF RESPONSIBLE INDIVIDUAL Professor Herbert C. Brown | | | 22b. TELEPHONE (Include Area Code) (317) 494-5316 | | 22c. OFFICE SYMBOL |

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

Unclassified

ABSTRACT

Enantioselective Ring-Cleavage of *meso*-Epoxides with B-Halodiisopinocampheylboranes.

N. N. Joshi, M. Srebnik, and Herbert C. Brown*

H. C. Brown and R. B. Wetherill Laboratories of Chemistry,
Purdue University, West Lafayette, Indiana 49707.

Cis-2-butene oxide
Abstract: B-Halodiisopinocampheylboranes, Ipc_2BX , in particular the bromide **1b**, and the Iodide **1c**, differentiate between the enantiotopic C-O bonds of *meso*-epoxides to furnish 1,2-halohydrins of moderate to excellent enantiomeric purity. Thus (-)-(1R, 2R)-2-bromocyclohexanol, (-)-(1R, 2R)-2-iodocyclohexanol, (-)-(1R, 2R)-2-iodocyclohex-4-en-1-ol and (-)-(1R, 2R)-2-bromocyclohex-4-en-1-ol are obtained in 84%, 91%, 63% and 95% ee respectively from the corresponding *meso*-epoxides and haloborane reagents. Simple recrystallization from pentane then gives halohydrins of essentially 100% ee; *cis*-2-butene oxide and *cis*-3-hexene oxide furnish the corresponding (1R, 2R)-iodohydrins in 78 and 69% ee respectively. In all cases the S carbon of the *meso*-epoxide is selectively cleaved. This is the first example of enantioselective ring cleavage of *meso*-epoxides to obtain optically active 1,2-halohydrins. *has* *6*

isomerized by 100%


| | |
|--------------------|--------------------------------------------|
| Accession For | |
| NTIS GRA&I | <input checked="checked" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By | |
| Distribution/ | |
| Availability Codes | |
| Avail and/or | |
| Dist | Special |
| A-1 | |

Enantioselective Ring-Cleavage of *meso*-Epoxides with B-Halodiisopinocampheylboranes.

N. N. Joshi, M. Srebnik, and Herbert C. Brown*

H. C. Brown and R. B. Wetherill Laboratories of Chemistry,
Purdue University, West Lafayette, Indiana 49707.

Asymmetric synthesis starting from *meso*-compounds is an increasingly important method for the preparation of optically active compounds. Various chemical¹ and enzymatic² procedures have been utilized for such transformations. Surprisingly, not much attention has been given to the enantioselective ring-cleavage of *meso*-epoxides which would lead to several classes of important compounds. We now report the synthesis of optically active 1,2-halohydrins using B-halodiisopinocampheylboranes, Ipc_2BX (**1a-c**).



1a X = Cl


1b X = Br

1c X = I

The cleavage of carbon-oxygen bonds with boron reagents, in particular with trihaloboranes, is a well documented and established procedure³. However, the high Lewis acidity of trihaloboranes and their tridentate nature can result in the unwanted rupture of bonds in compounds containing other sensitive functional groups. To increase the selectivity in carbon-oxygen bond cleavage, monohaloboranes in which two of the halogens on boron are replaced by alkyl⁴, alkoxy⁵, thioalkyl⁶, or nitrogen containing⁷ groups have recently been introduced. Such reagents indeed are selective and could by the proper choice of substituents cleave specific C-O bonds⁷. Recently Guindon and co-workers reported⁴ the stereospecific ring-

cleavage of various cyclic ethers with B-bromodimethylborane. Of particular consequence in the cleavage of epoxides with these type of reagents is the fact that the halohydrin products are consistent with a predominantly S_N2 type mechanism proposed for such reactions^{4,7,8}. This suggests that the asymmetric version should proceed with a high degree of enantiotopic differentiation of the C-O bonds of suitable *meso*-epoxides. Recently, the enantioselective cleavage of cyclohexene oxide with thiols and azides in the presence of zinc or copper tartrates has been successfully accomplished⁹. Except for this report, the cleavage of epoxides with chiral Lewis acids is an unexplored reaction. As part of our efforts in the field of asymmetric synthesis¹⁰, we are currently exploring reactions involving boron based chiral Lewis acids. The present communication describes our investigations on the enantioselective ring-opening of some *meso*-epoxides with B-halodiisopinocampheylboranes (Ipc_2BX)¹¹, both enantiomers of which are readily available from either (+)- or (-)- α -pinene.

First we examined the reaction between cyclohexene oxide and $d\text{Ipc}_2\text{BX}$ (the superscript "d" indicates that the reagent is derived from (+)- α -pinene) under various conditions, i.e., temperature, solvent and molarity. Due to the labile nature of the products, a non-oxidative work-up was developed (eq 1).




The course of the reaction was followed by ^{11}B -NMR¹². The reaction is very fast (<5 min) but proceeds with poor induction at 0°C . Not surprisingly, an inverse relationship between enantiomeric excess (% ee) and the reaction temperature was found to exist. It was established that Ipc_2BCl , Ipc_2BBr and Ipc_2BI require $-78^\circ\text{C}/3$ h, $-100^\circ\text{C}/2$ h and $-100^\circ\text{C}/0.5$ h respectively to furnish good enantioselection. Furthermore, the reaction is independent of solvent or molarity. Since **1b** and **1c** provided superior results (at least in

the case of cyclohexene oxide), we used these reagents subsequently for the preparation of optically active halohydrins (Table I).


In general [4.1.0]-oxaheptanes gave excellent initial optical yields of halohydrins¹³ We also found that by simple recrystallization from pentane, these could be upgraded to products of essentially 100 % ee. Cyclopentene oxide gave reduced chemical as well as optical yields. Simple acyclic epoxides were cleaved with optical inductions intermediate between those realised¹⁴ for the oxides of cyclohexene and cyclopentene. The % ee of the halohydrins (as their acetates) was determined using a chiral capillary GC column, Ni(HFB-1R-Cam)₂¹⁵.

The absolute configuration of the cyclic halohydrins was determined where possible by X-ray crystallography. The absolute configuration of the acyclic compounds was elucidated as follows (eq 2).




Dehalogenation with tributyltin hydride gave the known *secondary* alcohols¹⁶ which were analysed as their TPC-derivatives¹⁷ using capillary GC. Since the *meso*-epoxides are opened in an antiperiplanar manner, determination of the absolute configuration of one asymmetric center establishes the configuration of the other as well.

Of particular significance of this procedure is the fact that in addition to high optical induction¹⁸, the bond cleavage occurs in the same absolute sense to furnish (1R, 2R)-halohydrins in all cases examined, with ^dlpc₂BX¹⁹ These results are consistent with the cleavage occurring in an *anti* manner with inversion at the S carbon of the *meso*-epoxides (eq 3).



In addition, closure of the acyclic chiral non-racemic halohydrins yield the starting compounds, viz. a *cis*-epoxide, which would not be the case had the opening proceeded in a *syn* manner (eq 4).



In conclusion, the present methodology demonstrates the feasibility of cleaving *meso*-epoxides in an enantioselective manner either at the *R* or *S* carbon by selecting the appropriate chiral organoborane reagents. The transformation is general, providing highly valuable difunctionalized compounds in good to excellent enantiomeric purity from simple olefins. Synthesis of optically active halohydrins is but one of the many possible applications which we are currently exploring.

Acknowledgment: This work was supported in part by the Office of Naval Research. We thank Dr. Phil Fanwick for the X-ray crystallographic analysis.

References and Notes

- (1) (a) Fujita, E.; Nagao, Y.; Seno, K.; Takao, S.; Miyasaka, M.; Kimura, M.; Watson, W. H. *J. Chem. Soc. Perkin Trans. I* **1981**, 914. (b) Nagao, Y.; Ikeda, T.; Yagi, M.; Fujita, E.; Shiro, M. *J. Am. Chem. Soc.* **1982**, *104*, 2079. (c) Osakada, K.; Obana, M.; Ikariya, T.; Saburi, M.; Yoshikawa, S. *Tetrahedron Lett.* **1981**, *22*, 4297. (d) Fujisawa, T.; Watanabe, M.; Sato, T. *Chem. Lett.* **1984**, 2055.
- (2) Jakovac, I. J.; Goodbrand, H. B.; Lok, K. P.; Jones, J. B. *J. Am. Chem. Soc.* **1982**, *104*, 4659.
- (3) Bhat, M. V.; Kulkarni, S. U. *Synthesis* **1983**, 249.
- (4) Guindon, Y.; Therien, M.; Girard, Y.; Yoakim, C. *J. Org. Chem.* **1987**, *52*, 1680 and references cited therein.
- (5) Boekman, R. K. Jr.; Potenza, J. C. *Tetrahedron Lett.* **1985**, *26*, 1411.
- (6) (a) Williams, D. R.; Sakdarat, S. *ibid.* **1983**, *24*, 3965. (b) Corey, E. J.; Hua, D. H.; Seitz, S. P. *ibid.* **1984**, *25*, 3.
- (7) Bell, T. W.; Ciaccio, J. A. *ibid.* **1986**, *27*, 827.
- (8) Parkee, R. E.; Isaacs, N. S. *Chem. Rev.* **1959**, *59*, 737.
- (9) (a) Yamashita, H. *Chem. Lett.* **1987**, 525. (b) Yamashita, H.; Mukaiyama, T. *ibid.* **1985**, 1643.
- (10) For recent reviews see: (a) Brown, H. C.; Jadhav, P.K.; Singaram, B. In *Modern Synthetic Methods*; R. Scheffold, Ed.; Springer-Verlag: Berlin; **1986**, Vol. 4, p 307. (b) Srebnik, M.; Ramachandran, P. V. *Aldrichimica Acta* **1987**, *20*, 9.
- (11) Ipc_2BCl and Ipc_2BBr were prepared by passing HCl or HBr through a suspension of optically pure Ipc_2BH in pentane at 0°C . Ipc_2BI was made by reaction with I_2 . All the reagents were recrystallized and stored as stock solutions in pentane.
- (12) The reaction is quenched with pyridine which coordinates with the unreacted haloborane ($\delta = 18\text{-}20$ ppm) and not with the borinate ($\delta \approx 54$ ppm) formed during the course of the reaction.
- (13) **The preparation of (1R, 2R)-(-)-2-iodocyclohexanol:** A 100 ml flask protected with a positive pressure of nitrogen, was charged with Ipc_2BI (11 mmol, 0.25 M in pentane). The solution was cooled to -100°C and treated dropwise with cyclohexene oxide (10 mmol) dissolved in pentane (2 ml). After stirring for 1 h at 0°C , the

reaction was quenched by the addition of acetaldehyde (1.8 ml, excess). The mixture was allowed to warm gradually to room temperature and stirred for an additional 1 h. By that time ^{11}B -NMR indicated clean formation of boronate [$\text{R}^*\text{OB}(\text{OEt})\text{Ipc}$, $\delta = 31$ ppm]. The reaction mixture was diluted with pentane (50 ml) and treated with diethanolamine (12 mmol, 4 M in THF). After stirring for 30 min, the precipitated boronate-diethanolamine complex was filtered off, the filtrate was washed with water, brine, and dried over anhydrous Na_2SO_4 . The solution was concentrated under reduced pressure, and the residue purified by column chromatography followed by crystallization from pentane to provide the title compound (Table I, entry 3).

(14) No attempt was made to achieve optimum results.

(15) The column is available from: CC & CC (Capillary Column Complexation Chromatography), Postfach 14, D7402, Kirchentellinsfort, F.R.G. The results of this study will be published separately.






(16) Brown, H. C.; Ayyangar, N. R.; Zweifel, G. *J. Am. Chem. Soc.* **1964**, *80*, 397.

(17) Hoopes, E. A.; Peltzer, E. T.; Bada, J. L. *J. Chromat. Sci.* **1978**, *16*, 556.

(18) The origin of chirality in this transformation may be due to similar relative conformations of the two isopinocampheyl rings as was postulated for asymmetric hydroborations. See, Brown, H. C.; Jadhav, P. K.; Mandal, A. K. *Tetrahedron* **1981**, *37*, 3547.

(19) The use of $^{11}\text{Ipc}_2\text{BX}$ (derived from (-)- α -pinene) will obviously yield the corresponding (1S, 2S)-halohydrins.

Table 1. Optically Active 1,2-Halohydrins from *meso* - Epoxides and B-Halodiisopinocampheylboranes, ^d[pc₂BX^a

| entry | epoxide | 1,2-halohydrin | X | %yield ^b | bp °C/mmHg or mp °C | [α] _D ²³ D ₀ (c=5, CHCl ₃) (1R, 2R) | %ee ^c | abs. config. |
|-------|-----------------------------------|---------------------------------------------------------------------------------------|----|---------------------|------------------------|-------------------------------------------------------------------------------------|----------------------|--------------|
| 1 | cyclohexene oxide |  | Cl | 70 | 75-80/15 | | 22 | d |
| 2 | | | Br | 82 | 39-40 | -26.9 | 84(99) ^e | f |
| 3 | | | I | 89 | 44-45 | -31.6 | 91(100) ^e | g |
| 4 | 1,4-cyclohexadiene monoepoxide |  | Br | 72 | 90-95/15 | -42.0 | 63 | h |
| 5 | | | I | 75 | 37-38 | -141.8 | 95(100) ^e | g |
| 6 | cyclopentene oxide |  | I | 63 | 65-70/0.5 | -5.8 | 52 | c |
| 7 | <i>cis</i> -2-butene oxide |  | Br | 69 | 75-80/15 | -3.4 | 61 | h |
| 8 | | | I | 67 | 80-85/15 | -15.3 | 78 | h |
| 9 | <i>cis</i> -3-hexene oxide |  | Br | 71 | 90-95/15 | | 50 | h |
| 10 | | | I | 75 | 60-65/0.5 | +11.8 | 69 | h |

^aThe superscript "d" indicates that the reagent is derived from (+)-α-pinene. ^bIsolated by column chromatography followed by crystallization, or bulb-to-bulb distillation. ^cAnalysed as the corresponding acetates using Ni(HFB-1R-Cam)₂ capillary column. ^dBy analogy to the other examples. ^eOptical purity was upgraded by recrystallization from pentane. ^fBellucci, G.; Ingrosso, G.; Marioni, F.; Marsili, A.; Morelli, I. *Gazz. Chim Ital.* 1974, 104, 69. ^gDetermined by X-ray crystallography. ^hElucidated through the corresponding dehalogenated product. See text.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

| | <u>No. Copies</u> | | <u>No. Copies</u> |
|-----------------------------------------------------------------------------------------------------------------------------------|-----------------------|----------------------------------------------------------------------------------------------------------------|-----------------------|
| Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000 | 2 | Dr. David Young Code 334 NORDA NSTL, Mississippi 39529 | 1 |
| Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050 | 1 | Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555 | 1 |
| Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401 | 1 | Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380 | 1 |
| Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314 | 12 high quality | U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709 | 1 |
| DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401 | 1 | Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112 | 1 |
| Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000 | 1 | Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232 | 1 |

END

DATE

FILMED

DTIC

JULY 88